




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61F 2/02		A1	(11) International Publication Number: WO 99/21508
			(43) International Publication Date: 6 May 1999 (06.05.99)
(21) International Application Number: PCT/US98/22929		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 28 October 1998 (28.10.98)			
(30) Priority Data:		Published <i>With international search report.</i>	
08/959,061	28 October 1997 (28.10.97)	US	
08/958,571	28 October 1997 (28.10.97)	US	
09/181,071	27 October 1998 (27.10.98)	US	
(71) Applicant: VIVUS, INC. [US/US]; 605 East Fairchild Drive, Mountain View, CA 94043 (US).			
(72) Inventors: SMITH, William, L.; 6404 Wood Drive, Oakland, CA 94611 (US). DOHERTY, Paul, C., Jr.; 10461 Castine Avenue, Cupertino, CA 95014 (US). PLACE, Virgil, A.; P.O. Box 44555, 10 Ala Kahua, Kawaihae, HI 96743 (US).			
(74) Agents: REED, Dianne, E. et al.; Reed & Associates, 3282 Alpine Road, Portola Valley, CA 94028 (US).			

(54) Title: ADMINISTRATION OF ACTIVE AGENTS, INCLUDING 5-HT RECEPTOR AGONISTS AND ANTAGONISTS, TO TREAT PREMATURE EJACULATION



(57) Abstract

A method is provided for delaying the onset of ejaculation in an individual. The method preferably involves administration of an antidepressant drug, a serotonin agonist or antagonist, an adrenergic agonist or antagonist, an adrenergic neurone blocker, or a derivative analog thereof, within the context of an effective dosing regimen. The preferred mode of administration is transurethral; however, the selected active agent may also be delivered via intracavernosal injection or using alternative routes. Pharmaceutical formulations and kits are provided as well.

BEST AVAILABLE COPY

SOURCEONESM

For questions regarding this delivery, please contact Customer Services at 800-3-DIALOG.
Please have your DIALOG user number and order number available.

THE DIALOG CORPORATIONTM



11000 Regency Parkway, Suite 10, Cary NC. 27511 - 919-462-8600

This is not an invoice. Charges listed are for informational purposes only and do not include applicable tax, copyright or outside purchase fees or telecommunication charges. Copyrighted material reproduced with permission of copyright owner and further reproduction is prohibited or otherwise subject to restriction.

Document ID: WO9921508A1

For:

PPD, INC.
ATTN: BETSY CRITTENDEN
LEGAL
3151 SOUTH 17TH STREET
WILMINGTON
NC 28412
USA

Subject: DIALOG SOURCEONE
Order Date: 2000/06/26
Number: 200062
User Number: 153275
Subaccount: DIALOG SOURCEONE
Customer notes:

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SE	Sweden		
DK	Denmark	LK	Sri Lanka	SG	Singapore		
EE	Estonia	LR	Liberia				

ADMINISTRATION OF ACTIVE AGENTS, INCLUDING 5-HT RECEPTOR AGONISTS AND ANTAGONISTS, TO TREAT PREMATURE EJACULATION

5

TECHNICAL FIELD

This invention relates generally to methods and pharmaceutical compositions for treating sexual dysfunction; more particularly, the invention relates to treatment of premature ejaculation, such as by administration of compounds that are 5-HT (serotonin) receptor agonists and antagonists.

10

BACKGROUND ART

Premature ejaculation is a debilitating sexual dysfunction. This dysfunction can lead to an inability to enter or sustain relationships and can cause psychological damage to sufferers. Premature ejaculation can also impair reproductive success.

15

Previous methods of treating premature ejaculation include psychological therapies, topical anesthetics and the use of devices (U.S. Patent Nos. 5,535,758, 5,063,915, 5,327,910, and 5,468,212). All of these methods have significant drawbacks. Psychological therapies benefit only a subset of patients and require specialized therapists who may not be available to all patients, particularly in remote areas. Furthermore, psychological therapies cannot alleviate premature ejaculation resulting from non-psychological causes. Anesthetic agents decrease sensitivity of tissues, thereby diminishing sexual pleasure. Also, topical anesthetics can be transferred to sexual partners and thereby decrease their sensitivity and pleasure as well. With regard to devices, these can be awkward, inconvenient and embarrassing to use. Devices are highly conspicuous, and reveal the very condition which the suffering partner may prefer to conceal. Additionally, devices can cause irritation to one or both partners.

20

25

Methods for treating premature ejaculation by systemic administration of several different antidepressant compounds have been described (U.S. Patent Nos. 4,507,323, 4,940,731, 5,151,448, and 5,276,042; PCT Publication No. WO95/13072).

However, these drugs may not be effective for all patients, and the side effects of these drugs can halt treatment or impair patient compliance. Disease states or adverse interactions with other drugs may contraindicate the use of these compounds or require lower dosages that may not be effective to delay the onset of ejaculation. Additionally, the stigma of mental
5 illness associated with antidepressant therapy can discourage patients from beginning or continuing such treatments.

Administration of the antidepressant fluoxetine has been claimed to treat premature ejaculation (U.S. Patent No. 5,151,448). However, the administration of fluoxetine has many undesired aspects. Patients with hepatic or renal impairments may not
10 be able to use fluoxetine due to its metabolism in the liver and excretion via the kidney. Systemic events during fluoxetine treatment involving the lungs, kidneys or liver have occurred, and death has occurred from overdoses. In addition, side effects of oral fluoxetine administration include hair loss, nausea, vomiting, dyspepsia, diarrhea, anorexia, anxiety, nervousness, insomnia, drowsiness, fatigue, headache, tremor, dizziness, convulsions,
15 sweating, pruritis, and skin rashes. Fluoxetine interacts with a range of drugs, often by impairing their metabolism by the liver.

U.S. Patent No. 4,940,731 describes the oral or parenteral administration of sertraline for treating premature ejaculation. It has been recognized that sertraline shares many of the same problems as fluoxetine; see Martindale, *The Extra Pharmacopoeia*, 31st
20 edition, at p. 333 (London: The Royal Pharmaceutical Society, 1996). Sertraline is metabolized in the liver, and is excreted in the urine and feces. Thus, patients with cirrhosis must take lower doses, and caution must be exercised when administering sertraline to patients with renal impairment. Individuals taking monoamine oxidase inhibitors cannot take sertraline due to the risk of toxicity, leading to memory changes, confusion, irritability,
25 chills, pyrexia and muscle rigidity. Side effects resulting from oral sertraline administration include nausea, diarrhea, dyspepsia, insomnia, somnolence, sweating, dry mouth, tremor and mania. Rare instances of coma, convulsions, fecal incontinence and gynecomastia have occurred in patients undergoing sertraline therapy.

U.S. Patent No. 5,276,042 describes the administration of paroxetine for the
30 treatment of premature ejaculation. Paroxetine is predominantly excreted in the urine, and

-3-

decreased doses are recommended in patients with hepatic and renal impairments. Like sertraline, paroxetine cannot be given to patients undergoing treatment with a monoamine oxidase inhibitor. Side effects from oral administration of paroxetine include hyponatremia, asthenia, sweating, nausea, decreased appetite, oropharynx disorder, somnolence, dizziness, insomnia, tremor, anxiety, impaired micturition, weakness and paresthesia.

Thus there is a need for a method of treating premature ejaculation that requires no specialized psychological therapy, can be used conveniently and without embarrassment, and does not involve the problems associated with prior therapeutic methods.

Serotonin, or 3-(β -aminoethyl)-5-hydroxyindole (5-hydroxytryptophan, or "5-HT") is a neurotransmitter in the central nervous system which is known to play an important role in the pathogenesis of affective illness. Several different 5-HT receptor types have been identified, including 5-HT₁, 5-HT₂ and 5-HT₃, which are further divided into a number of different subtypes, e.g., 5-HT_{1A}, 5-HT_{1B}, 5-HT_{1D}, 5-HT_{1E} and 5-HT_{1F}. It has now been discovered that administration of various serotonin agonists and antagonists is quite effective in the treatment of premature ejaculation, and addresses a number of the above-noted deficiencies in the art. Accordingly, in a preferred embodiment, the present invention is directed to the administration of serotonin agonists and antagonists, preferably 5-HT₃ receptor antagonists (also referred to herein as "5-HT₃ antagonists") and 5-HT₄ receptor agonists (also referred to herein as "5-HT₄ agonists"), in the treatment of premature ejaculation. In alternative embodiments, the invention is also directed to the treatment of premature ejaculation using other types of active agents, including adrenergic neurone blockers and adrenergic agonists and antagonists.

DISCLOSURE OF INVENTION

Accordingly, it is a primary object of the invention to address the above-described need in the art by providing a novel method for treating premature ejaculation by administering an effective amount of a serotonin antagonist or agonist to an individual in need of such therapy.

It is another object of the invention to provide such a method wherein the pharmacologically active agent is administered orally.

It is a further object of the invention to provide such a method wherein the pharmacologically active agent is administered parenterally.

It is another object of the invention to provide such a method wherein the pharmacologically active agent is administered buccally.

5 It is also an object of the invention to provide such a method wherein the pharmacologically active agent is administered nasally.

It is another object of the invention to provide such a method wherein the pharmacologically active agent is administered transurethrally.

10 It is an additional object of the invention to provide such a method wherein the pharmacologically active agent is administered via intracavernosal injection.

It is another object of the invention to provide a novel method for treating premature ejaculation by locally administering an effective amount of a pharmacologically active agent to an individual in need of such therapy, wherein the active agent is an antidepressant drug, a serotonin antagonist or agonist (as above), an adrenergic neurone blocker, or an adrenergic agonist or antagonist.

It is yet a further object of the invention to provide pharmaceutical formulations for carrying out the aforementioned methods.

It is another object of the invention to provide a kit capable of use by an individual in carrying out the aforementioned methods.

20 Additional objects, advantages and novel features of the invention will be set forth in part in the description which follows, and in part will become apparent to those skilled in the art upon examination of the following, or may be learned by practice of the invention.

In a first aspect of the invention, a method is provided for treating premature ejaculation, the method comprising administering to an individual in need of such treatment a pharmaceutical formulation containing a serotonin antagonist or agonist. Administration of the pharmaceutical formulation is carried out within the context of a predetermined dosing regimen such that the agent is effective in the treatment of premature ejaculation. Drug delivery may be accomplished through any route effective to provide relief from

premature ejaculation, including oral, parenteral, buccal, rectal, topical, transdermal, transurethral, and intracavernosal injection.

In a further aspect of the invention, a method is provided for treating premature ejaculation, the method comprising locally administering to an individual in need of such treatment a pharmaceutical formulation containing a selected pharmacologically active agent, wherein, in this embodiment, the pharmacologically active agent is an antidepressant drug (preferably a selective serotonin reuptake inhibitor), a serotonin agonist or antagonist, an adrenergic neurone blocker, or an adrenergic agonist or antagonist. Administration of the pharmaceutical formulation is carried out within the context of a predetermined dosing regimen such that the agent is effective in the treatment of premature ejaculation. Drug delivery is preferably effected transurethrally, but may also be administered via intracavernosal injection or using topical or transdermal administration.

In another aspect of the invention, a pharmaceutical formulation is provided for carrying out the methods of the invention. The pharmaceutical formulation comprises an effective amount of an active agent as provided herein, a pharmacologically acceptable carrier or vehicle, and, optionally (i.e., in topical, transdermal or transurethral formulations), an enhancer. Other types of components may be incorporated into the formulation as well, e.g., excipients, surfactants, preservatives (e.g., antioxidants), stabilizers, enzyme inhibitors, chelating agents, and the like, as will be appreciated by those skilled in the art of pharmaceutical formulation preparation and drug delivery.

In another aspect of the invention, a kit is provided to assist an individual in administering a drug to treat premature ejaculation. Generally, the kit will include the following components: a pharmaceutical formulation comprising an active agent as provided herein; a device for effecting delivery of the pharmaceutical formulation; a container housing the pharmaceutical formulation during storage and prior to use; and instructions for carrying out drug administration in a manner effective to delay the onset of ejaculation.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 illustrates an embodiment of a transurethral therapeutic device which may be used in conjunction with the present methods.

5

MODES FOR CARRYING OUT THE INVENTION

Overview and Definitions:

Before describing the present invention in detail, it is to be understood that this invention is not limited to particular drugs or drug delivery systems, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

It must be noted that, as used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmacologically active agent" includes a combination of two or more pharmacologically active agents, reference to "a transurethral permeation enhancer" includes combinations of two or more enhancers, and the like.

In describing and claiming the present invention, the following terminology will be used in accordance with the definitions set out below.

The terms "active agent," "drug" and "pharmacologically active agent" are used interchangeably herein to refer to a chemical material or compound which, when administered to an organism (human or animal) induces a desired pharmacologic effect. Included are derivatives and analogs of those compounds or classes of compounds specifically mentioned which also induce the desired pharmacologic effect.

The terms "transurethral," "intraurethral" and "urethral" to specify the preferred mode of administration herein are used interchangeably to refer to delivery of the drug into the urethra such that the drug contacts and passes through the wall of the urethra. As noted elsewhere herein, the transurethral administration preferably involves delivery of the drug at least about 3 cm and more preferably at least about 7 cm into the urethra.

-7-

The term "intracavernosal" as used herein refers to an alternative mode of drug administration and involves injection into one or both corpora of the corpora cavernosal tissues of the penis.

By the term "transdermal" delivery, applicants intend to include both
5 transdermal (or "percutaneous") and transmucosal administration, i.e., delivery by passage of a drug through the skin or mucosal tissue and into the bloodstream. The term "body surface" will sometimes be used herein to refer to either the skin or the mucosal tissue. "Transdermal" delivery is also intended to encompass passage through scrotal skin.

The term "topical administration" is used in its conventional sense to mean
10 delivery of a topical drug or pharmacologically active agent to the skin or mucosa.

The term "local administration" as used herein generally refers to transurethral delivery, topical administration, and administration by intracavernosal injection.

"Penetration enhancement" or "permeation enhancement" as used herein relates to an increase in the permeability of the skin, mucosa or urethral membrane to the selected
15 pharmacologically active agent, i.e., so that the rate at which the drug permeates therethrough is increased.

"Carriers" or "vehicles" as used herein refer to carrier materials suitable for drug administration. Carriers and vehicles useful herein include any such materials known in the art, e.g., any liquid, gel, solvent, liquid diluent, solubilizer, or the like, which is nontoxic and
20 which does not interact with other components of the composition in a deleterious manner.

The term "premature ejaculation" as used herein intends a sexual dysfunction wherein a male is unable to control the ejaculatory process to a degree sufficient to satisfy a partner. Generally, "premature ejaculation" refers to persistent or recurring ejaculation with minimal stimulation before or during sexual intercourse. The term includes both
25 "congenital" or "lifelong" premature ejaculation and "primary" or "acquired" premature ejaculation as set forth, for example, in U.S. Patent No. 5,151,448, and in Male Infertility and Sexual Dysfunction at p. 386 (New York: Springer-Verlag, 1997). See also Diagnostic and Statistical Manual of Mental Disorders (Washington, D.C.: American Psychiatric Association, 1994).

By an "effective" amount of a drug or pharmacologically active agent is meant a nontoxic but sufficient amount of the drug or agent to provide the desired effect.

Active Agents for Treating Premature Ejaculation:

5 In order to carry out the method of the invention, a selected pharmacologically active agent is administered to an individual with a history of premature ejaculation. In a first embodiment, the active agent may be administered orally, parenterally, buccally, rectally, transdermally, or locally by intracavernosal injection, topical administration, or delivery to the urethra. In this first embodiment, suitable pharmacologically active agents
10 that can be administered to treat premature ejaculation include, but are not limited to:

 serotonin agonists including 2-methyl serotonin, buspirone, ipsaperone, tiaspirone, gepirone, D-lysergic acid diethylamide ("LSD"), ergot alkaloids, 8-hydroxy-(2-N,N-dipropyl-amino)-tetraline, 1-(4-bromo-2,5-dimethoxyphenyl)-2-aminopropane, cisapride, sumatriptan, *m*-chlorophenylpiperazine, trazodone, zacopride and mezacopride;
15 and

 serotonin antagonists including ondansetron, granisetron, metoclopramide, tropisetron, dolasetron, palonosetron, trimethobenzamide, methysergide, risperidone, ketanserin, ritanserin, clozapine, amitriptyline, MDL 100,907 (R(+)- α -(2,3-dimethoxyphenyl)-1-[2-(4-fluorophenyl)ethyl]-4-piperidine-methanol) (Marion Merrell
20 Dow), azatadine, cyproheptadine, fenclonine, chlorpromazine and mianserin.

 Serotonin, or 5-HT, plays a variety of roles throughout the body, regulating smooth muscle and platelet function, and acting as a neurotransmitter in the central nervous system. Serotonin acts through specific cellular receptors which comprise at least 15 individual subtypes. These receptors can be grouped into at least four families based on
25 structural and functional characteristics. The 5-HT₁, 5-HT₂ and 5-HT₄ families are G-protein coupled receptors linked to enzymatic and electrical effector systems, while the 5-HT₃ receptor is a gated ion channel. Serotonin agonists are agents which mimic the effect of serotonin on at least one serotonin receptor subtype. For example, 5-HT₄ agonists mimic the effect of serotonin on at least one 5-HT₄ receptor. Conversely, serotonin antagonists are

agents which block the effect of serotonin on at least one serotonin receptor subtype; 5-HT₃ antagonists block the effect of serotonin on the 5-HT₃ receptor.

Preferred active agents are 5-HT₃ antagonists and 5-HT₄ agonists. 5-HT₃ receptors can be found, for example, on parasympathetic terminals in the gastrointestinal tract and in the central nervous system, both of which participate in the emetic response. 5-HT₄ receptors are found throughout the body, including on nerve terminals in the CNS, the gastrointestinal tract, and on smooth muscle and secretory cells. 5-HT₄ receptors activate adenylyl cyclase, and are involved in the regulation of secretion and peristalsis. Examples of 5-HT₃ antagonists include ondansetron, ergot alkaloids, granisetron, metoclopramide, trimethobenzamide, tropisetron, dolasetron, batanopride and zacopride. Examples of 5-HT₄ agonists include cisapride and D-lysergic acid diethylamide.

In another embodiment of the invention, an active agent as set forth below is administered locally to treat premature ejaculation. In this embodiment, suitable pharmacologically active agents include, but are not limited to:

antidepressant drugs including amesergide, amineptine, amitriptyline, amoxapine, benactyzine, brofaromine, bupropion, butriptyline, cianopramine, citalopram, clomipramine, clorgyline, clovoxamine, demexiptiline, desipramine, dibenzepin, dimetacrine, dothiepin, doxepin, etoperidone, femoxetine, fezolamine, fluoxetine, fluvoxamine, ifoxetine, imipramine, iprindole, isocarboxazid, levoprotiline, lofepramine, maprotiline, medifoxamine, melitracen, metapramine, methylphenidate, mianserin, milnacipran, minaprine, mirtazapine, moclobemide, nefazodone, nialamide, nomifensine, nortriptyline, opipramol, oxaflozane, oxaprotiline, oxitriptan, paroxetine, phenelzine, pirlindole, propizepine, protriptyline, quinupramine, rolipram, selegiline, sertraline, setiptiline, sibutramine, teniloxazine, tianeptine, tofenacin, toloxatone, tranylcypromine, trazodone, trimipramine, tryptophan, venlafaxine, viloxazine, viqualine and zimeldine; serotonin agonists including 2-methyl serotonin, buspirone, ipsaperone, tiaspirone, gepirone, LSD; ergot alkaloids, 8-hydroxy-(2-N,N-dipropylamino)-tetraline, 1-(4 bromo 2,5 dimethoxyphenyl)-2-aminopropane, cisapride, sumatriptan, m-chlorophenylpiperazine, trazodone, zacopride and mezacopride;

serotonin antagonists including ondansetron, granisetron, metoclopramide, tropisetron, dolasetron, palonosetron, trimethobenzamide, methysergide, risperidone, ketanserin, ritanserin, clozapine, amitriptyline, MDI 100,907, azatadine, cyproheptadine, fenclonine, chlorpromazine, mianserin, zacopride and mezacopride;

5 adrenergic agonists including methoxamine, methpentermine, metaraminol, mitodrine, clonidine, apraclonidine, guanfacine, guanabenz, methyl dopa, amphetamine, methamphetamine, epinephrine, norepinephrine, ethylnorepinephrine, phenylephrine, ephedrine, pseudoephedrine, methylphenidate, pemoline, naphazoline, tetrahydrozoline, oxymetazoline, xylometazoline, phenylpropanolamine, phenylethylamine, dopamine,
10 dobutamine, colterol, isoproterenol, isotharine, metaproterenol, terbutaline, metaraminol, tyramine, hydroxyamphetamine, ritodrine, prenalterol, albuterol, isoetharine, pirbuterol, bitolterol, fenoterol, formoterol, procaterol, salmeterol, mephenterine and propylhexedrine;

 adrenergic neurone blockers including bethanidine, debrisoquine, guabenxan, guanadrel, guanazodine, guanethidine, guanoclor and guanoxan; and

15 adrenergic antagonists including phenoxybenzamine, phentolamine, tolazoline, prazosin, terazosin, doxazosin, trimazosin, yohimbine, ergot alkaloids, labetalol, ketanserin, urapidil, alfuzosin, bunazosin, tamsulosin, chlorpromazine, haloperidol, phenothiazines, butyrophenones, propranolol, nadolol, timolol, pindolol, metoprolol, atenolol, esmolol, acebutolol, bopindolol, carteolol, oxprenolol, penbutolol, carvedilol, medroxalol, naftopidil,
20 bucindolol, levobunolol, metipranolol, bisoprolol, nebivolol, betaxolol, carteolol, celiprolol, sotalol, propafenone and indoramin.

 Some agents, as may be seen, are encompassed by more than one of the above categories, e.g., serotonin antagonists and antidepressants, or serotonin agonists and antagonists.

25 The active agent may be administered in the form of a pharmaceutically acceptable salt, ester, amide or prodrug or combination thereof. However, conversion of inactive ester, amide or prodrug forms to an active form must occur prior to or upon reaching the target tissue or cell. Salts, esters, amides and prodrugs of the active agents may be prepared using standard procedures known to those skilled in the art of synthetic organic
30 chemistry and described, for example, by J. March, Advanced Organic Chemistry:

Reactions, Mechanisms and Structure, 4th Ed. (New York: Wiley-Interscience, 1992). For example, acid addition salts are prepared from the free base (typically wherein the neutral form of the drug has a neutral $-NH_2$ group) using conventional means, involving reaction with a suitable acid. Generally, the base form of the drug is dissolved in a polar organic solvent such as methanol or ethanol and the acid is added thereto. The resulting salt either precipitates or may be brought out of solution by addition of a less polar solvent. Suitable acids for preparing acid addition salts include both organic acids, e.g., acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, malic acid, malonic acid, succinic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like, as well as inorganic acids, e.g., hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. An acid addition salt may be reconverted to the free base by treatment with a suitable base. Conversely, preparation of basic salts of acid moieties which may be present on a drug are prepared in a similar manner using a pharmaceutically acceptable base such as sodium hydroxide, potassium hydroxide, ammonium hydroxide, calcium hydroxide, trimethylamine, or the like. Preparation of esters involves functionalization of hydroxyl and/or carboxyl groups which may be present within the molecular structure of the drug. The esters are typically acyl-substituted derivatives of free alcohol groups, i.e., moieties which are derived from carboxylic acids of the formula $RCOOH$ where R is alkyl, and preferably is lower alkyl. Esters can be reconverted to the free acids, if desired, by using conventional hydrogenolysis or hydrolysis procedures. Preparation of amides and prodrugs can be carried out in an analogous manner.

Other derivatives and analogs of the active agents may be prepared using standard techniques known to those skilled in the art of synthetic organic chemistry, or may be deduced by reference to the pertinent literature. In addition, chiral active agents may be in isomerically pure form, or they may be administered as a racemic mixture of isomers.

Pharmaceutical Formulations and Modes of Administration:

Depending on the intended mode of administration, the pharmaceutical compositions may be in the form of solid, semi-solid or liquid dosage forms, such as, for example, tablets, suppositories, pills, capsules, powders, liquids, suspensions, creams, ointments, lotions or the like, preferably in unit dosage form suitable for single administration of a precise dosage. The compositions will include an effective amount of the selected drug in combination with a pharmaceutically acceptable carrier and, in addition, may include other pharmaceutical agents, adjuvants, diluents, buffers, etc. The amount of active compound administered will, of course, be dependent on the subject being treated, the subject's weight, the manner of administration and the judgment of the prescribing physician.

For solid compositions, conventional nontoxic solid carriers include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talc, cellulose, glucose, sucrose, magnesium carbonate, and the like. Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, etc., an active compound as described herein and optional pharmaceutical adjuvants in an excipient, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to thereby form a solution or suspension. If desired, the pharmaceutical composition to be administered may also contain minor amounts of nontoxic auxiliary substances such as wetting or emulsifying agents, pH buffering agents and the like, for example, sodium acetate, sorbitan monolaurate, triethanolamine sodium acetate, triethanolamine oleate, etc. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington's Pharmaceutical Sciences, referenced above.

For oral administration, the composition will generally take the form of a tablet or capsule, or may be an aqueous or nonaqueous solution, suspension or syrup. Tablets and capsules are preferred oral administration forms. Tablets and capsules for oral use will generally include one or more commonly used carriers such as lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. When liquid suspensions are used, the active agent may be combined with emulsifying and suspending

-13-

agents. If desired, flavoring, coloring and/or sweetening agents may be added as well. Other optional components for incorporation into an oral formulation herein include, but are not limited to, preservatives, suspending agents, thickening agents, and the like.

Parenteral administration, if used, is generally characterized by injection.

- 5 Injectable formulations can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solubilization or suspension in liquid prior to injection, or as emulsions. Preferably, sterile injectable suspensions are formulated according to techniques known in the art using suitable carriers, dispersing or wetting agents and suspending agents. The sterile injectable formulation may also be a sterile injectable
10 solution or a suspension in a nontoxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils, fatty esters or polyols are conventionally employed as solvents or suspending media. A more recently revised approach for parenteral administration involves use of a slow release or sustained release
15 system, such that a constant level of dosage is maintained. See, e.g., U.S. Patent No. 3,710,795.

- Intracavernosal injection can be carried out by use of a syringe any other suitable device. An example of a hypodermic syringe useful herein, that can be used for simultaneous injection into both corpora, is described in U.S. Patent No. 4,127,118 to
20 Latorre. The injection is made on the dorsum of the penis by placement of the needle to the side of each dorsal vein and inserting it deep into the corpora.

- The active agent can be administered in a pharmaceutical formulation suitable for transurethral drug delivery. The formulation contains one or more selected carriers or excipients, such as water, silicone, waxes, petroleum jelly, polyethylene glycol ("PEG"),
25 propylene glycol ("PG"), liposomes, sugars such as mannitol and lactose, and/or a variety of other materials, with polyethylene glycol and derivatives thereof particularly preferred.

- Depending on the drug administered, it may be desirable to incorporate a transurethral permeation enhancer in the urethral dosage form. Examples of suitable transurethral permeation enhancers include dimethylsulfoxide ("DMSO"), dimethyl
30 formamide ("DMF"), N,N-dimethylacetamide ("DMA"), decylmethylsulfoxide

5 (“C₁₀MSO”), polyethylene glycol monolaurate (“PEGML”), glycerol monolaurate, lecithin, the 1-substituted azacycloheptan-2-ones, particularly 1-*n*-dodecylcyclazacycloheptan-2-one (available under the trademark Azone® from Nelson Research & Development Co., Irvine, CA), SEPA® (available from Macrochem Co., Lexington, MA), alcohols (e.g., ethanol),
detergents (such as Tergitol®, Nonoxynol-9® and TWEEN-80®) and the like.

Transurethral formulations may additionally include one or more enzyme inhibitors effective to inhibit drug-degrading enzymes which may be present in the urethra. Such enzyme inhibiting compounds may be determined by those skilled in the art by reference to the pertinent literature and/or using routine experimental methods. Additional
10 optional components include excipients, preservatives (e.g., antioxidants), chelating agents, solubilizing agents (e.g., surfactants), and the like, as will be appreciated by those skilled in the art of drug formulation preparation and delivery.

Transurethral drug administration, as explained in PCT Publication WO 91/16021 and in U.S. Patent Nos. 5,242,391, 5,474,535, 5,686,093 and 5,773,020 to Place et al., can be carried out in a number of different ways using a variety of urethral dosage forms.
15 For example, the drug can be introduced into the urethra from a flexible tube, squeeze bottle, pump or aerosol spray. The drug may also be contained in coatings, pellets or suppositories which are absorbed, melted or bioeroded in the urethra. In certain embodiments, the drug is included in a coating on the exterior surface of a penile insert. A preferred drug delivery
20 device for administering a drug transurethally is shown in Figure 1. It is preferred, although not essential, that the drug be delivered at least about 3 cm into the urethra, and preferably at least about 7 cm into the urethra. Generally, delivery at about 3 cm to about 8 cm into the urethra will provide effective results in conjunction with the present method.

Urethral suppository formulations containing PEG or a PEG derivative are
25 particularly preferred urethral dosage forms herein, and may be conveniently formulated using conventional techniques, e.g., compression molding, heat molding or the like, as will be appreciated by those skilled in the art and as described in the pertinent literature and pharmaceutical texts. See, for example, Remington: The Science and Practice of Pharmacy, 19th Ed. (Easton, PA: Mack Publishing Co., 1995), which discloses typical methods of
30 preparing pharmaceutical compositions in the form of urethral suppositories. The PEG or

PEG derivative preferably has a molecular weight M_w in the range of about 200 to 2500, more preferably in the range of about 1000 to 2000. Suitable polyethylene glycol derivatives include polyethylene glycol fatty acid esters, for example, polyethylene glycol monostearate, polyethylene glycol sorbitan esters, e.g., polysorbates, and the like. It is also
5 preferred that urethral suppositories contain one or more solubilizing agents effective to increase the solubility of the active agent in the PEG or other transurethral vehicle.

The solubilizing agent may be a nonionic, anionic, cationic or amphoteric surfactant. Nonionic surfactants include: long-chain fatty acids, i.e., acids having the structural formula $\text{CH}_3(\text{CH}_2)_m\text{COOH}$ where m is an integer in the range of 8 to 16; fatty
10 alcohols, that is, alcohols having the structural formula $\text{CH}_3(\text{CH}_2)_m\text{C(H)OH}$, such as lauryl, cetyl and stearyl alcohols; glyceryl esters such as the naturally occurring mono-, di- and triglycerides; and esters of fatty alcohols or other alcohols such as propylene glycol, polyethylene glycol, sorbitan, sucrose, and cholesterol. Examples of water-soluble nonionic surfactant derivatives include sorbitan fatty acid esters (such as those sold under the
15 tradename Span®), polyoxyethylene sorbitan fatty acid esters (such as those sold under the tradename TWEEN®), polyoxyethylene fatty acid esters (such as those sold under the tradename Myrj®), polyoxyethylene steroidal esters, polyoxypropylene sorbitan fatty acid esters, polyoxypropylene fatty acid esters, polyoxypropylene steroidal esters, polyoxyethylene ethers (such as those sold under the tradename Brij®), polyglycol ethers
20 (such as those sold under the tradename Tergitol®), and the like. Preferred nonionic surfactants for use as the solubilizing agent herein are polyglycol ether, polyoxyethylene sorbitan trioleate, sorbitan monopalmitate, polysorbate 80, polyoxyethylene 4-lauryl ether, propylene glycol, and mixtures thereof. Anionic surfactants which may be used as the solubilizing agent herein include long-chain alkyl sulfonates, carboxylates, and sulfates, as
25 well as alkyl aryl sulfonates, and the like. Preferred anionic surfactants are sodium dodecyl sulfate, dialkyl sodium sulfosuccinate (e.g., sodium bis-(2-ethylhexyl)-sulfosuccinate), sodium 7-ethyl-2-methyl-4-dodecyl sulfate and sodium dodecylbenzene sulfonate. Cationic surfactants which may be used to solubilize the active agent are generally long-chain amino salts or quaternary ammonium salts, e.g., decyltrimethylammonium bromide,
30 dodecyltrimethylammonium bromide, tetradecyltrimethylammonium bromide,

tetradecyltrimethylammonium chloride, and the like. Amphoteric surfactants are generally, although not necessarily, compounds which include a carboxylate or phosphate group as the anion and an amino or quaternary ammonium moiety as the cation. These include, for example, various polypeptides, proteins, alkyl betaines, and natural phospholipids such as lecithins and cephalins. Other suitable solubilizing agents include glycerin, propylene glycol, fatty acids and fatty alcohols. The solubilizing agent will be present in the range of approximately 0.01 wt.% to 40 wt.%, more preferably in the range of approximately 5.0 wt.% to 40 wt.%, and most preferably in the range of approximately 10.0 wt.% to 40 wt.%.

It may be desirable to deliver the active agent in a urethral dosage form which provides for controlled or sustained release of the agent. In such a case, the dosage form typically comprises a biocompatible, biodegradable material, typically a biodegradable polymer. Examples of such polymers include polyester, polyalkylcyanoacrylate, polyorthoester, polyanhydride, albumin, gelatin and starch. As explained, for example, in PCT Publication No. WO96/40054, these and other polymers can be used to provide biodegradable microparticles which enable controlled and sustained drug release, in turn minimizing the required dosing frequency.

The urethral suppository will preferably, although not necessarily, be on the order of 2 to 20 mm, preferably 5 to 10 mm in length and less than about 5 mm, preferably less than about 2 mm in width. The weight of the suppository form will typically be in the range of approximately 1 mg to 50 mg. However, it will be appreciated by those skilled in the art that the size of the suppository can and will vary, depending on the potency of the drug, the nature of the formulation, and other factors.

In Figure 1, a suitable transurethral drug delivery device is shown generally at 10. The device comprises a transurethral inserter 11 having an easily graspable segment 12 that has opposing symmetrically concave surfaces 13 and 14 adapted to be held by two fingers. Drug is contained within a urethral suppository (not shown) within shaft 15, which is sized to fit within the urethra. A longitudinal plunger, the tip of which is seen at 16, is slidably insertable into the longitudinal bore contained within shaft 15. To extrude drug into the urethra, shaft 15 is inserted into the urethra, and plunger tip 16 is pushed into segment 12. The inserter 11 is then removed. Prior to use, and during storage, the device is capped

-17-

with elongate cap 17 which fits snugly over flange 18 at the proximal end of shaft 15. The cap 17 is provided with a series of parallel ridges 19 to facilitate gripping of the cap and removal from inserter 11.

Although the transurethral drug delivery device shown in Figure 1 represents a preferred device for use herein, again, it should be emphasized that a wide variety of device configurations and urethral dosage forms can be used.

Examples of other devices suited to deliver a drug transurethrally are those described and illustrated in PCT Publication No. WO 91/16021 and in U.S. Patent Nos. 5,242,391, 5,474,535, 5,686,093 and 5,773,020 to Place et al.

The devices can either be manufactured under sterile conditions, thereby eliminating the need for post-manufacturing sterilization, or they can be manufactured under non-sterile conditions and then subsequently sterilized by any suitable technique, e.g., radiation sterilization. The devices can be manufactured by typical plastic forming and coating processes known in the art, including molding extrusion, heat forming, dip coating, and the like.

Transurethral drug delivery may involve an "active" delivery mechanism such as iontophoresis, electroporation or phonophoresis. Devices and methods for delivering drugs in this way are well known in the art. Iontophoretically assisted drug delivery is, for example, described in PCT Publication No. WO96/40054, cited above. Briefly, the active agent is driven through the urethral wall by means of an electric current passed from an external electrode to a second electrode contained within or affixed to a urethral probe.

The compounds of the invention may also be delivered through the skin or mucosal tissue using conventional transdermal drug delivery systems, i.e., transdermal "patches" wherein the agent is typically contained within a laminated structure that serves as a drug delivery device to be affixed to the body surface. In such a structure, the drug composition is typically contained in a layer, or "reservoir," underlying an upper backing layer. The laminated device may contain a single reservoir, or it may contain multiple reservoirs. In one embodiment, the reservoir comprises a polymeric matrix of a pharmaceutically acceptable contact adhesive material that serves to affix the system to the skin during drug delivery. Examples of suitable skin contact adhesive materials include, but

are not limited to, polyethylenes, polysiloxanes, polyisobutylenes, polyacrylates, polyurethanes, and the like. Alternatively, the drug-containing reservoir and skin contact adhesive are present as separate and distinct layers, with the adhesive underlying the reservoir which, in this case, may be either a polymeric matrix as described above, or it may
5 be a liquid or gel reservoir, or may take some other form. The backing layer in these laminates, which serves as the upper surface of the device, functions as the primary structural element of the laminated structure and provides the device with much of its flexibility. The material selected for the backing layer should be substantially impermeable to the active agent and any other materials that are present.

10 Alternatively, the pharmaceutical compositions of the invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

15 The pharmaceutical compositions of the invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, propellants such as fluorocarbons or nitrogen, and/or other conventional
20 solubilizing or dispersing agents.

Preferred formulations for topical drug delivery are ointments and creams. Ointments are semisolid preparations which are typically based on petrolatum or other petroleum derivatives. Creams containing the selected active agent, are, as known in the art, viscous liquid or semisolid emulsions, either oil-in-water or water-in-oil. Cream bases are
25 water-washable, and contain an oil phase, an emulsifier and an aqueous phase. The oil phase, also sometimes called the "internal" phase, is generally comprised of petrolatum and a fatty alcohol such as cetyl or stearyl alcohol; the aqueous phase usually, although not necessarily, exceeds the oil phase in volume, and generally contains a humectant. The emulsifier in a cream formulation is generally a nonionic, anionic, cationic or amphoteric
30 surfactant. The specific ointment or cream base to be used, as will be appreciated by those

skilled in the art, is one that will provide for optimum drug delivery. As with other carriers or vehicles, an ointment base should be inert, stable, nonirritating and nonsensitizing.

Formulations for buccal administration include tablets, lozenges, gels and the like. Alternatively, buccal administration can be effected using a transmucosal delivery system as known to those skilled in the art.

The pharmaceutical formulation may contain one or more pharmacologically active agents in addition to the serotonin agonist or antagonist. Vasoactive agents, particularly vasodilators, are preferred additional agents.

Suitable vasoactive agents include, but are not limited to: nitrates and like compounds such as nitroglycerin, isosorbide dinitrate, erythrityl tetranitrate, amyl nitrate, sodium nitroprusside, molsidomine, linsidomine chlorhydrate ("SIN-1"), S-nitroso-N-acetyld,l-penicillamine ("SNAP"), S-nitroso-N-cysteine and S-nitroso-N-glutathione ("SNO-GLU") and diazenium diolates ("NONOates"); long and short acting α -blockers such as phenoxybenzamine, dibenamine, doxazosin, terazosin, phentolamine, tolazoline, prazosin, trimazosin, alfuzosin, tamsulosin and indoramin; ergot alkaloids such as ergotamine and ergotamine analogs, e.g., acetergamine, brazergoline, bromerguride, cianergoline, delorgotril, disulergine, ergonovine maleate, ergotamine tartrate, etisulergine, lergotril, lysergide, mesulergine, metergoline, metergotamine, nicergoline, pergolide, propisergide, proterguride and terguride; antihypertensive agents such as diazoxide, hydralazine and minoxidil; vasodilators such as nimodipine, pinacidil, cyclandelate, dipyrindamole and isoxsuprine; chlorpromazine; haloperidol; yohimbine; Rec15/2739; trazodone; naturally occurring prostaglandins such as PGE₀, PGE₁, PGA₁, PGB₁, PGF_{1 α} , 19-hydroxy-PGA₁, 19-hydroxy-PGB₁, PGE₂, PGA₂, PGB₂, 19-hydroxy-PGA₂, 19-hydroxy-PGB₂, PGE₃, PGF_{3 α} ; semisynthetic or synthetic derivatives of natural prostaglandins, including carboprost tromethamine, dinoprost tromethamine, dinoprostone, lipoprost, gemeprost, metenoprost, suiprostone and tiaprost; and vasoactive intestinal peptides. Prazosin, prostaglandin E₀, prostaglandin E₁ and prostaglandin E₂ are particularly preferred vasoactive agents to be co-administered with the active agent.

The amount of active agent administered, and the dosing regimen used, will, of course, be dependent on the particular drug selected, the age and general condition of the

-20-

subject being treated, the severity of the subject's condition, and the judgment of the prescribing physician. Generally, the daily dosage when administered locally will be less than the dosage normally given in conjunction with systemic modes of administration, and typically, the drug will be administered one to four times daily or, with some active agents, just prior to intercourse. Alternatively, a large initial loading dose can be used to achieve effective levels of the agent and can be followed by smaller doses to maintain those levels. A typical daily dose of an active agent as administered locally is generally in the range of approximately 0.1 to 500 mg. Depending on the half-life of the drug and the availability via the chosen route of administration, the dosing regimen can be modulated in order to achieve satisfactory control of the onset of ejaculation.

Kits:

The invention also encompasses a kit for patients to carry out the present method of treating premature ejaculation. The kit contains the pharmaceutical formulation to be administered, a device for administering the formulation (e.g., a transurethral drug delivery device such as shown in Figure 1, or a syringe); a container, preferably sealed, for housing the drug and device during storage and prior to use, and instructions for carrying out drug administration in an effective manner. The formulation may consist of the drug in unit dosage form. The kit may contain multiple formulations of different dosages of the same agent. The kit may also contain multiple formulations of different active agents. The instructions may be in written or pictograph form, or can be on recorded media including audio tape, video tape, or the like.

Use in Conjunction with Venous Flow Control ("VFC") Device:

In an alternative embodiment of the invention, the pharmacologically active agent is administered in combination with a venous flow control device such as that described in PCT Publication No. WO 97/47260, entitled "Venous Flow Control Element for Maintaining Penile Erection." Preferred devices are formed from a length of flexible tubing having an integral fastening means, so as to provide for readily adjustable venous flow control when applied to the penis. The device is applied to the base of the penis prior

-21-

to and during sexual intercourse, such that it effectively enhances retention of blood within the penis without substantially obstructing arterial inflow or becoming too constrictive during the erectile process. Use of the VFC device also enables enhanced effectiveness of local drug therapy, in that the active agent is retained within the penis, allowing movement
5 into the corpus cavernosa. This produces smooth muscle response and a consistent erectile response. In this embodiment, a kit will include the venous flow control device in addition to the components noted above, along with instructions for using the device.

10

It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof, that the foregoing description as well as the examples which follow are intended to illustrate and not limit the scope of the invention. Other aspects, advantages and modifications within the scope of the invention will be
15 apparent to those skilled in the art to which the invention pertains.

EXAMPLE 1

A pharmaceutical formulation for transurethral administration is prepared by mixing 0.1-25 mg of the serotonin agonist sumatriptan with polyethylene glycol, molecular
20 weight (M_w) approximately 4000, and heating the mixture to a temperature just high enough to produce a sumatriptan-polymer melt. The sumatriptan-glycol mixture can then be poured into a mold suitable to provide a sumatriptan suppository approximately 5 mm in length and 1.5 mm in width, and allowed to cool. The suppository so provided is a unit dosage form suitable for transurethral administration. If desired, the sumatriptan-glycol mixture may be
25 allowed to cool on the tip of a rod adapted to be inserted into the urethra.

-22-

EXAMPLE 2

The procedure of Example 1 is repeated, except that 1-50 mg of the serotonin agonist cisapride is substituted for sumatriptan. A suppository suitable for transurethral administration of a unit dosage of cisapride is thus provided.

5

EXAMPLE 3

The procedure of Example 1 is repeated, except that 0.1-25 mg of the serotonin antagonist risperidone is substituted for sumatriptan. A suppository suitable for transurethral administration of a unit dosage of risperidone is thus provided.

10

EXAMPLE 4

The procedure of Example 1 is repeated, except that 1-100 mg of the serotonin antagonist clozapine is substituted for sumatriptan. A suppository suitable for transurethral administration of a unit dosage of clozapine is thus provided.

15

EXAMPLE 5

The procedures of the foregoing Examples are repeated, except that cocoa butter is substituted for polyethylene glycol.

20

EXAMPLE 6

A penile insert coated with a serotonin agonist or antagonist is prepared as follows. An ethylene vinyl acetate (28% VA) rod is formed into an insert having a shaft approximately 10 cm long with a spherical, blunted tip. A dipping bath comprising a 50-50 weight blend of PEG 1450 and PEG 4000 and sufficient agent to attain the desired concentration in the coating is prepared and heated to 70°C. The insert is suspended by its head, dipped into the dipping bath and removed. A penile insert suitable for transurethral administration of agent is thus provided.

30

EXAMPLE 7

A pharmaceutical formulation suitable for injection can be made by dissolving 4 mg of the 5-HT₃ antagonist ondansetron hydrochloride dihydrate, 9.0 mg of sodium chloride, 0.5 mg of citric acid monohydrate and 0.25 mg of sodium citrate dihydrate in water
5 to give a 2 mL solution.

EXAMPLE 8

A pharmaceutical formulation suitable for oral administration can be prepared by admixing either 4 or 8 mg of the 5-HT₃ antagonist ondansetron hydrochloride dihydrate
10 with appropriate amounts of lactose, microcrystalline cellulose, pregelatinized starch, hydroxypropyl methylcellulose, magnesium stearate and titanium dioxide and forming the mixture into tablets.

EXAMPLE 9

A pharmaceutical formulation suitable for oral administration can be made
15 suspending the 5-HT₄ agonist cisapride monohydrate at 1 mg/mL in an aqueous solution containing appropriate amounts of sodium chloride, hydroxypropyl methylcellulose, methylparaben, microcrystalline cellulose and carboxymethylcellulose sodium, polysorbate
20, propylparaben, and sorbitol.

20

EXAMPLE 10

A pharmaceutical formulation suitable for oral administration can be prepared by admixing either 10 or 20 mg of the 5-HT₄ agonist cisapride monohydrate with appropriate amounts of lactose monohydrate, microcrystalline cellulose, polysorbate 20, povidone, corn starch, magnesium stearate and colloidal silicon dioxide and forming the
25 mixture into tablets.

EXAMPLE 11

A pharmaceutical formulation containing the 5-HT₃ antagonist ondansetron hydrochloride dihydrate for transdermal administration is prepared by mixing prazosin
30 hydrochloride, a vehicle and an adhesive, and casting the solution onto the backing layer. If

desired, a skin permeation enhancer may be included. This release liner is then sealed to the backing liner, and a protective layer is applied to the release liner. The device is employed by removing the protective layer and applying the release liner to the skin.

5

EXAMPLE 12

A pharmaceutical formulation containing the 5-HT₃ antagonist ondansetron hydrochloride dihydrate for topical application is prepared by mixing 25 to 500 mg prazosin hydrochloride in 100 mL of propylene glycol. In addition, antioxidants such as mixed tocopherols may be added in the range of from about 0.1 mg to 5 mg. Viscosity modifiers
10 such as paraffin wax, lanolin wax or other compatible waxes may also be added to the formulation. The formulation is applied locally for the treatment of premature ejaculation.

EXAMPLE 13

The patients for the study are drawn from a waiting list for psychosexual
15 therapy of a sexology outpatient department and through advertisement. The inclusion criteria are that the patients be heterosexual; be aged 18-75 years, be involved in a sexual relationship with a female partner during the previous 6 months, the partner is able to participate in the study, and the patient experience premature ejaculation.

Patients are excluded if they used psychoactive medication, are receiving any
20 therapy for sexual dysfunction, have inhibited male orgasm, are alcohol or substance abusers, score greater than 14 on the 24-item Hamilton Rating Scale for Depression (HAM-D) indicating clinically significant depression, and for other clinically significant medical disease or symptomatology. For the duration of the study, the patients are required to abstain from alcohol.

25 Patients are randomly assigned to double-blind treatment with an antidepressant, a serotonin agonist or antagonist, or a derivative or analog thereof, or a placebo. Patients who remain in the study for at least 3 weeks are included in the statistical analysis. Patients are provided with formulations prepared in the preceding Examples for transurethral administration or matching placebo. During the study, the patients do not use
30 condoms or topical anesthetics.

Patients and their partners are assessed at the end of each week. Efficacy measures include (1) frequency of attempted intercourse, (2) latency to attempted ejaculation (from penetration), (3) frequency of successful attempts at intercourse without premature ejaculation, (4) number of incidences of premature ejaculation, and (5) time to ejaculation as reported by both the patients and their partners based on subjective measurements. In addition, the patients are assessed using the following psychopathological rating instruments: Hamilton Depression Rating Scale (HDRS), Clinical Global Impression (CGI), COVI Anxiety Scale, Atypical Depressive Disorders Scale-Changes (ADDS-C) and an adverse event form.

10 The response to treatment is scored. The mean or median values are analyzed by using parametric tests including Friedman two-way analysis of variance (ANOVA) and the Wilcoxon matched-pairs signed-ranks test to assess differences in measurements within the groups, and the Mann-Whitney test for differences between the groups. Differences between groups on discrete variables are tested for statistical significance by using Fisher's exact test. A two-tailed p value ≤ 0.05 is considered significant for the analysis.

15 Each of the formulations prepared and administered is expected to be effective in treating premature ejaculation. The use of the formulations results in an extension of intravaginal ejaculation latency time in patients with premature ejaculation. The data from patient-rated and partner-rated latency to ejaculation is compared. Both of these parameters show greater improvement when formulations of the preceding Examples are administered compared with the placebo group. The administration of the drugs by intracavernosal injection, transdermal or topical application is expected to yield similar results.

EXAMPLE 14

25 The procedure of Example 13 is repeated, except that a flexible, adjustable venous flow control device is used prior to and during sexual intercourse, in combination with the drug therapy described. Substantially the same results are expected.

-26-

EXAMPLE 15

The procedure of Example 13 is repeated, except that the drug is administered orally. Substantially the same results are expected.

5

EXAMPLE 16

The procedure of Example 13 is repeated, except that the drug is administered transdermally. Substantially the same results are expected.

EXAMPLE 17

10 The procedure of Example 13 is repeated, except that the drug is administered buccally. Substantially the same results are expected.

EXAMPLE 18

15 A pharmaceutical formulation containing an α_1 -adrenergic antagonist for transurethral administration is prepared by mixing 0.25 to 5 mg prazosin hydrochloride with a suitable amount of polyethylene glycol, typically in the range of approximately 1-5 g and having a molecular weight (M_w) of approximately 2000, and heating the mixture to a temperature just high enough to produce a drug-polymer melt. The mixture can then be poured into a mold suitable to provide a unit dosage form suitable for transurethral
20 administration. This procedure can be used with various drugs, PEGs, and additional components, e.g., enhancers or the like, to prepare pharmaceutical formulations suitable for urethral administration in the treatment of premature ejaculation.

EXAMPLE 19

25 A pharmaceutical formulation containing an adrenergic blocking agent for transurethral administration is prepared by mixing 1-10 mg terazosin with polyethylene glycol, molecular weight (M_w) approximately 4000, and heating the mixture to a temperature just high enough to produce a terazosin-polymer melt. The terazosin-glycol mixture can then be poured into a mold suitable to provide a terazosin suppository approximately 5 mm in length
30 and 1.5 mm in width, and allowed to cool, and allowed to cool. The suppository so

provided is a unit dosage form suitable for transurethral administration. If desired, the terazosin-glycol mixture may be allowed to cool on the tip of a rod adapted to be inserted into the urethra.

5

EXAMPLE 20

The procedure of Example 19 is repeated, except that 1-10 mg doxazosin is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of the adrenergic antagonist is thus provided.

10

EXAMPLE 21

The procedure of Example 19 is repeated, except that 1-100 mg of the antidepressant clomipramine is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of clomipramine is thus provided.

15

EXAMPLE 22

The procedure of Example 19 is repeated, except that 1-100 mg of the antidepressant sertraline is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of sertraline is thus provided.

20

EXAMPLE 23

The procedure of Example 19 is repeated, except that 0.1-25 mg of the serotonin agonist sumatriptan is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of sumatriptan is thus provided.

25

EXAMPLE 24

The procedure of Example 19 is repeated, except that 1-50 mg of the serotonin agonist cisapride is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of cisapride is thus provided.

30

EXAMPLE 25

The procedure of Example 19 is repeated, except that 0.1-25 mg of the serotonin antagonist risperidone is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of risperidone is thus provided.

5

EXAMPLE 26

The procedure of Example 19 is repeated, except that 1-100 mg of the serotonin antagonist clozapine is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of clozapine is thus provided.

10

EXAMPLE 27

The procedure of Example 19 is repeated, except that 1-100 mg of the adrenergic neurone blocker guanadrel is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of guanadrel is thus provided.

15

EXAMPLE 28

The procedure of Example 19 is repeated, except that 0.1-50 mg of the adrenergic agonist albuterol is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of albuterol is thus provided.

20

EXAMPLE 29

The procedure of Example 19 is repeated, except that 1-20 mg of the adrenergic agonist methoxamine is substituted for terazosin. A suppository suitable for transurethral administration of a unit dosage of methoxamine is thus provided.

25

EXAMPLE 30

The procedures of the foregoing Examples are repeated, except that cocoa butter is substituted for polyethylene glycol.

EXAMPLE 31

A pharmaceutical formulation containing an adrenergic antagonist for urethral administration is prepared by dissolving terazosin in a sufficient quantity of water to equal a predetermined percentage of the final suppository weight. Glycerin (70%) is then added,
5 followed by addition of Pharmagel A or B (20%). Suppositories are then prepared as described in Example 1.

EXAMPLE 32

The procedure of Example 31 is repeated, except that doxazosin is substituted
10 for terazosin. A suppository suitable for transurethral administration of a unit dosage of the adrenergic antagonist is thus provided.

EXAMPLE 33

The procedure of Example 18 is repeated, except that a flexible, adjustable
15 venous flow control device is used prior to and during sexual intercourse, on combination with the drug therapy described. Substantially the same results are expected.

EXAMPLE 34

A pharmaceutical formulation containing an α_1 -adrenergic antagonist for
20 intracavernosal injection is prepared by mixing prazosin hydrochloride in sterile isotonic buffer to provide a unit dosage in the range of approximately 0.25 to 5 mg. Patients receive a single injection of the sterile aqueous formulation containing prazosin hydrochloride, using a 26-30 gp needle connected to a syringe. Intracavernous pressure is measured by inserting a separate 21-25 gp plastic needle connected to a pressure transducer into the
25 opposite cavernous body.

EXAMPLE 35

A pharmaceutical formulation containing an α_1 -adrenergic antagonist for
transdermal administration is prepared by mixing prazosin hydrochloride, a vehicle and an
30 adhesive, and casting the solution onto the backing layer. If desired, a skin permeation

-30-

enhancer may be included. This release liner is then sealed to the backing liner, and a protective layer is applied to the release liner. The device is employed by removing the protective layer and applying the release liner to the skin.

5

EXAMPLE 36

A pharmaceutical formulation containing an α_1 -adrenergic antagonist for topical application is prepared by mixing 25 to 500 mg prazosin hydrochloride in 100 mL of propylene glycol. In addition, antioxidants such as mixed tocopherols may be added in the range of from about 0.1 mg to 5 mg. Viscosity modifiers such as paraffin wax, lanolin wax or other compatible waxes may also be added to the formulation. The formulation is applied locally for the treatment of premature ejaculation.

10

EXAMPLE 37

The patients for the study are drawn from a waiting list for psychosexual therapy of a sexology outpatient department and through advertisement. The inclusion criteria are that the patients be heterosexual, be aged 18-75 years; be involved in a sexual relationship with a female partner during the previous 6 months, the partner is able to participate in the study, and the patient experience premature ejaculation.

15

Patients are excluded if they use psychoactive medication, are receiving any therapy for sexual dysfunction, are alcohol or substance abusers, have inhibited male orgasm, score greater than 14 on the 24-item Hamilton Rating Scale for Depression (HAM-D) indicating clinically significant depression, and for other clinically significant medical disease or symptomatology. For the duration of the study, the patients are required to abstain from alcohol.

20

Patients are randomly assigned to double-blind treatment with an antidepressant, a serotonin agonist or antagonist, an adrenergic agonist or antagonist, an adrenergic neurone blocker, or a derivative or analog thereof, or a placebo. Patients who remain in the study for at least 3 weeks are included in the statistical analysis. Patients are provided with formulations prepared in the preceding Examples for transurethral

25

-31-

administration or matching placebo. During the study, the patients do not use condoms or topical anesthetics.

Patients and their partners are assessed at the end of each week. Efficacy measures include (1) frequency of attempted intercourse, (2) latency to attempted ejaculation (from penetration), (3) frequency of successful attempts at intercourse without premature ejaculation, (4) number of incidences of premature ejaculation, and (5) time to ejaculation as reported by both the patients and their partners based on subjective measurements. In addition, the patients are assessed using the following psychopathological rating instruments: Hamilton Depression Rating Scale (HDRS), Clinical Global Impression (CGI), COVI Anxiety Scale, Atypical Depressive Disorders Scale-Changes (ADDS-C) and an adverse event form.

The response to treatment is scored. The mean or median values are analyzed by using parametric tests including Friedman two-way analysis of variance (ANOVA) and the Wilcoxon matched-pairs signed-ranks test to assess differences in measurements within the groups, and the Mann-Whitney test for differences between the groups. Differences between groups on discrete variables are tested for statistical significance by using Fisher's exact test. A two-tailed p value ≤ 0.05 is considered significant for these analysis.

Each of the formulations prepared and administered is expected to be effective in treating premature ejaculation. The use of the formulations results in an extension of intravaginal ejaculation latency time in patients with premature ejaculation. The data from patient-rated and partner-rated latency to ejaculation is compared. Both of these parameters show greater improvement when formulations of the preceding Examples are administered compared with the placebo group. The administration of the drugs by intracavernosal injection, transdermal or topical application is expected to yield similar results.

25

EXAMPLE 38

The procedure of Example 37 is repeated, except that a flexible, adjustable venous flow control device is used prior to and during sexual intercourse, in combination with the drug therapy described. Substantially the same results are expected.

30

EXAMPLE 39

The procedure of Example 37 is repeated, except that the drug is administered transdermally. Substantially the same results are expected.

CLAIMS

1. A method for delaying the onset of ejaculation in a human individual, comprising administering to the individual an effective amount of a pharmaceutical formulation containing an active agent comprising a serotonin agonist, a serotonin antagonist, or a combination thereof, wherein the active agent is effective to delay the onset of ejaculation by the individual during sexual intercourse.
2. The method of claim 1, wherein the formulation is administered transurethally.
3. The method of claim 1, wherein the formulation is administered by intracavernosal injection.
4. The method of claim 1, wherein the formulation is administered transdermally.
5. The method of claim 1, wherein the formulation is administered topically.
6. The method of claim 1, wherein the formulation is administered orally.
7. The method of claim 1, wherein the formulation is administered parenterally.
8. The method of claim 1, wherein the formulation is administered buccally.
9. The method of claim 1, wherein the formulation is administered nasally.
10. The method of claim 1, wherein the agent is a serotonin agonist.
11. The method of claim 10, wherein the serotonin agonist is a 5-HT₄ agonist.

-34-

12. The method of claim 1, wherein the agent is a serotonin antagonist.

13. The method of claim 12, wherein the serotonin antagonist is a 5-HT₃ antagonist.

5

14. The method of claim 1, wherein a predetermined dose of the agent is administered to the individual one to four times in a twenty-four hour period.

15. A pharmaceutical formulation for delaying the onset of ejaculation in a male individual, comprising a urethral dosage form of an active agent comprising a serotonin agonist, a serotonin antagonist, or a combination thereof, and a pharmaceutically acceptable carrier or excipient.

10

16. The formulation of claim 15, wherein the agent is a serotonin agonist.

15

17. The formulation of claim 16, wherein the serotonin agonist is a 5-HT₄ agonist.

18. The formulation of claim 15, wherein the agent is a serotonin antagonist.

20

19. The formulation of claim 18, wherein the serotonin antagonist is a 5-HT₃ antagonist.

20. A method for delaying the onset of ejaculation in a male individual, comprising locally administering to the individual an effective amount of a pharmaceutical formulation containing a pharmacologically active agent of a type and in an amount effective to delay the onset of ejaculation by the individual during sexual intercourse, wherein the agent is selected from the group consisting of antidepressants, serotonin agonists, serotonin antagonists, adrenergic agonists, adrenergic antagonists, adrenergic neurone blockers, and derivatives and analogs thereof.

25

30

-35-

21. A kit for delaying the onset of ejaculation in an individual, comprising: a pharmacologically active agent selected from the group consisting of antidepressants, serotonin agonists, serotonin antagonists, adrenergic agonists, adrenergic antagonists, adrenergic neurone blockers and combinations thereof; a drug delivery means for
5 administering the agent; a container for housing the agent and drug delivery means; and instructions for using the agent and drug delivery means to administer the drug within the context of a dosing regimen effective to treat premature ejaculation.


10

15

20

25

30



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/22929

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61F 2/02

US CL : 424/423

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/423

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,587,167 A (CHOI et al) 24 December 1996, see Abstract, claims.	1, 5
X	US 5,276,042 A (CRENSHAW et al.) 04 January 1994, see Abstract and col. 3, lines 4-5.	1, 4, 8, 9
X	WO 91/16021 A1 (ALZA CORPORATION) 31 October 1991, see Abstract; page 5, lines 22-32; page 6, lines 30-31.	1, 2, 7, 20, 21

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	* T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* A* document defining the general state of the art which is not considered to be of particular relevance	* X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* B* earlier document published on or after the international filing date	* Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* Z* document member of the same patent family
* O* document referring to an oral disclosure, use, exhibition or other means	
* P* document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

05 DECEMBER 1998

Date of mailing of the international search report

14 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CARLOS AZPURU

Telephone No. (703) 308-1235

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.